arXiv:1703.03203v1 [math.CA] 9 Mar 2017

# Error estimates for Riemann sums of some singular functions

Pavel Gurevich*, Sergey Tikhomirov†

March 10, 2017

**Abstract**

In this short note, we obtain error estimates for Riemann sums of some singular functions.

Let $N \in \mathbb{N}$, and let $z_0 = -1$ or $z_0 = 0$. For a function $F_1 : [z_0, 1] \to \mathbb{R}$ denote the error estimate of the Riemann sum of the integral $\int_{z_0}^1 F_1(x)dx$ by

$$R_n := \int_{z_0}^1 F_1(x)dx - \left( \frac{1}{2n} F_1(z_0) + \frac{1}{2n} F_1(1) + \sum_{k=z_0 n+1}^{n-1} \frac{1}{n} F_1\left(\frac{k}{n}\right) \right).$$

We prove the following 3 propositions, which are used, e.g., in [1].

**Proposition 1.** *Assume that a function $F_1(x)$ can be represented as*

$$F_1(x) = c_1(1-x)^{1/2} + c_2(1-x)^{3/2} + \tilde{F}_1(x),$$

*where $c_1, c_2 \in \mathbb{R}$ and $\tilde{F}_1 \in C^2[z_0, 1]$. Then*

1. *There exists $L_1 = L_1(F_1, N) > 0$ such that*

$$|R_n| \le L_1 \frac{1}{n^{3/2}}, \quad n \ge N.$$

2. *If, additionally, $\tilde{F}_1(x) = c_3(1-x)^{5/2} + c_4(1-x)^{7/2} + \bar{F}_1(x)$, where $c_3, c_4 \in \mathbb{R}$ and $\bar{F} \in C^4[z_0, 1]$, then there exists $\bar{L}_1 = \bar{L}_1(F_1, N) > 0$ such that*

$$\left|(n+1)^2 R_{n+1} - n^2 R_n\right| \le \bar{L}_1 \frac{1}{n^{1/2}} \quad n \ge N.$$

*Free University of Berlin, RUDN University, email: gurevich@math.fu-berlin.de
†Saint-Petersburg State Univeristy; email: s.tikhomirov@spbu.ru

**Proposition 2.** *Assume that a function $F_2(x)$ can be represented as*

$$F_2(x) = c_1(1-x)^{-1/2} + c_2(1-x)^{1/2} + \tilde{F}_2(x), \tag{1}$$

*where $c_1 > 0$, $c_2 \in \mathbb{R}$, and $\tilde{F}_2 \in C^1[z_0, 1]$. Then there exists $L_2 = L_2(F_2, N) > 0$, $L_2^* = L_2^*(F_2, N) > c_1$ and $l_2 = l_2(F_2, N)$ such that*

$$L_2^* \frac{1}{n^{1/2}} - l_2 \frac{1}{n} \le \int_{z_0}^{1} F_2(x)dx - \sum_{k=z_0 n}^{n-1} \frac{1}{n} F_2\left(\frac{k}{n}\right) \le L_2 \frac{1}{n^{1/2}}, \quad n \ge N. \tag{2}$$

*In particular, $\left| \sum\limits_{k=z_0 n}^{n-1} \frac{1}{n} F_2\left(\frac{k}{n}\right) \right|$ is bounded.*

**Proposition 3.** *Assume that $F_3 \in C^0[-1, 1)$ and $|(1-x)^{3/2} F_3(x)|$ is bounded on $[-1, 1)$. Then there exists $L_3 = L_3(F_3, N) > 0$ such that*

$$\left| \sum_{|k| \le n-1} \frac{1}{n} F_3\left(\frac{k}{n}\right) \right| \le L_3 n^{1/2}, \quad n \ge N.$$

Here we provide the proofs of item 2 of Proposition 1 and Proposition 2 for the case $z_0 = -1$. Item 1 of Proposition 1 and the case $z_0 = 0$ can be proved similarly. Proposition 3 is straightforward.

## Proof of item 2 of Proposition 1

Consider the function $\Phi(x) := F_1(x) + F_1(-x)$. Then

$$n^2 R_n = n^2 I_{F1} - \left( \frac{n}{2}\Phi(0) + \frac{n}{2}\Phi(1) + \sum_{k=1}^{n-1} n\Phi\left(\frac{k}{n}\right) \right),$$

where $I_{F1} := \int_{-1}^{1} F_1(x)dx$. Therefore,

$$\begin{aligned}(n+1)^2 R_{n+1} - n^2 R_n &= (2n+1) I_{F1} - \left( \frac{\Phi(0) + \Phi(1)}{2} + \sum_{k=1}^{n} (n+1)\Phi\left(\frac{k}{n+1}\right) - \sum_{k=1}^{n-1} n\Phi\left(\frac{k}{n}\right) \right) \\ &= \Sigma_{1,n} + \Sigma_{2,n}, \end{aligned} \tag{3}$$

where

$$\begin{aligned} \Sigma_{1,n} &:= (n+1) I_{F1} - \left( \frac{\Phi(0) + \Phi(1)}{2} + \sum_{k=1}^{n} \Phi\left(\frac{k}{n+1}\right) \right), \\ \Sigma_{2,n} &:= n(I_{F1} - \Phi(0) - \Sigma_{\Phi,n}), \quad \Sigma_{\Phi,n} := \sum_{k=1}^{n} \left( \Phi\left(\frac{k}{n+1}\right) - \Phi\left(\frac{k-1}{n}\right) \right). \end{aligned}$$

Note that $\Sigma_{1,n} = (n+1)R_{n+1}$. Hence, Proposition 1 (item 1) implies that, for some $L_1 > 0$,

$$|\Sigma_{1,n}| \le L_1 \frac{1}{\sqrt{n+1}}. \tag{4}$$

Estimates of $\Sigma_{2,n}$ are more subtle. First, note that $\frac{k}{n+1} - \frac{k-1}{n} = \frac{n+1-k}{n(n+1)}$ and, hence,

$$\Phi\left(\frac{k}{n+1}\right) - \Phi\left(\frac{k-1}{n}\right) = \frac{n+1-k}{n(n+1)}\Phi'\left(\frac{k}{n+1}\right) - \frac{1}{2}\left(\frac{n+1-k}{n(n+1)}\right)^2 \Phi''\left(\frac{k}{n+1}\right) + \frac{1}{6}\left(\frac{n+1-k}{n(n+1)}\right)^3 \Phi'''(\xi_k),$$

where $\xi_k \in \left[\frac{k-1}{n}, \frac{k}{n+1}\right]$. Therefore,

$$\Sigma_{\Phi,n} = \frac{n+1}{n}\sum_{k=1}^{n} \frac{1}{n+1}\frac{n+1-k}{n+1}\Phi'\left(\frac{k}{n+1}\right) - \frac{1}{2}\frac{n+1}{n^2}\sum_{k=1}^{n}\frac{1}{n+1}\left(\frac{n+1-k}{n+1}\right)^2 \Phi''\left(\frac{k}{n+1}\right) + \frac{1}{6}\frac{n+1}{n^3}\sum_{k=1}^{n}\frac{1}{n+1}\left(\frac{n+1-k}{n+1}\right)^3 \Phi'''(\xi_k)\,. \tag{5}$$

Set

$$\Phi_1(x) := (1-x)\Phi'(x), \quad \Phi_2(x) := (1-x)^2\Phi''(x), \quad \Phi_3(x) := (1-x)^3\Phi'''(x),$$

$$I_{\Phi 1} := \int_0^1 \Phi_1(x)dx, \quad I_{\Phi 2} := \int_0^1 \Phi_2(x)dx, \tag{6}$$

$$\Sigma_{\Phi 1,n} := I_{\Phi 1} - \sum_{k=1}^{n}\frac{1}{n+1}\frac{n+1-k}{n+1}\Phi'\left(\frac{k}{n+1}\right) - \frac{1}{2(n+1)}\Phi'(0), \tag{7}$$

$$\Sigma_{\Phi 2,n} := I_{\Phi 2} - \sum_{k=1}^{n}\frac{1}{n+1}\left(\frac{n+1-k}{n+1}\right)^2 \Phi''\left(\frac{k}{n+1}\right) - \frac{1}{2(n+1)}\Phi''(0), \tag{8}$$

$$\Sigma_{\Phi 3,n} := \sum_{k=1}^{n}\frac{1}{n+1}\left(\frac{n+1-k}{n+1}\right)^3 \Phi'''(\xi_k)\,. \tag{9}$$

Since $\xi_k \le \dfrac{k}{n+1}$, we have

$$\left|\left(\frac{n+1-k}{n+1}\right)^3 \Phi'''(\xi_k)\right| \le |\Phi_3(\xi_k)|. \tag{10}$$

Proposition 1 (item 1) and (10) imply that

$$|\Sigma_{\Phi 1,n}| \le K_{\Phi 1}\frac{1}{(n+1)^{3/2}}, \quad |\Sigma_{\Phi 2,n}| \le K_{\Phi 2}\frac{1}{(n+1)^{3/2}}, \quad |\Sigma_{\Phi 3,n}| \le K_{\Phi 3}, \tag{11}$$

$$K_{\Phi 1} := L_1(\Phi_1, N), \quad K_{\Phi 2} := L_1(\Phi_2, N), \quad K_{\Phi 3} := \sup_{x\in(0,1)} |\Phi_3(x)|,$$

In this notation, (5) takes the form

$$\Sigma_{\Phi,n} = \frac{n+1}{n}\left(I_{\Phi 1} - \frac{\Phi'(0)}{2(n+1)} - \Sigma_{\Phi 1,n}\right) - \frac{n+1}{2n^2}\left(I_{\Phi 2} - \frac{\Phi''(0)}{2(n+1)} - \Sigma_{\Phi 2,n}\right) + \frac{n+1}{6n^3}\Sigma_{\Phi 3,n}.$$

Finally, taking into account (4), we conclude that

$$\begin{aligned}\Sigma_{2,n} = nI_{F1} - (n+1)I_{\Phi 1} + \frac{1}{2}\frac{n+1}{n}I_{\Phi 2} - \left(n\Phi(0) - \frac{1}{2}\Phi'(0) + \frac{1}{4n}\Phi''(0)\right)\\ + \left((n+1)\Sigma_{\Phi 1,n} - \frac{1}{2}\frac{n+1}{n}\Sigma_{\Phi 2,n} - \frac{1}{6}\frac{n+1}{n^2}\Sigma_{\Phi 3,n}\right).\end{aligned}$$

Using the relations $I_{F1} = I_{\Phi 1} + \Phi(0)$ and $I_{\Phi 2} = 2I_{\Phi 1} - \Phi'(0)$, we obtain

$$\begin{aligned}\Sigma_{2,n} &= nI_{\Phi 1} + n\Phi(0) - (n+1)I_{\Phi 1} + \frac{n+1}{n}I_{\Phi 1} - \frac{1}{2}\frac{n+1}{n}\Phi'(0)\\ &\quad - \left(n\Phi(0) - \frac{1}{2}\Phi'(0) + \frac{1}{4n}\Phi''(0)\right) + \left((n+1)\Sigma_{\Phi 1,n} - \frac{1}{2}\frac{n+1}{n}\Sigma_{\Phi 2,n} - \frac{1}{6}\frac{n+1}{n^2}\Sigma_{\Phi 3,n}\right)\\ &= \frac{1}{n}I_{\Phi 1} - \frac{1}{4n}\Phi''(0) - \frac{1}{2n}\Phi'(0) + \left((n+1)\Sigma_{\Phi 1,n} - \frac{1}{2}\frac{n+1}{n}\Sigma_{\Phi 2,n} - \frac{1}{6}\frac{n+1}{n^2}\Sigma_{\Phi 3,n}\right).\end{aligned}$$

The latter equality and estimates (11) yield

$$\begin{aligned}|\Sigma_{2,n}| &\le \frac{1}{n}\left(|I_{\Phi 1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi 3}}{3}\right) + K_{\Phi 1}\frac{1}{n^{1/2}} + \frac{K_{\Phi 2}}{2}\frac{1}{n^{3/2}}\\ &\le \frac{1}{n^{1/2}}\left(K_{\Phi 1} + \left(|I_{\Phi 1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi 3}}{3}\right)\frac{1}{N^{1/2}} + \frac{K_{\Phi 2}}{2}\frac{1}{N}\right).\end{aligned}$$

Combining this with (3) and (4), we have

$$|(n+1)^2 R_{n+1} - n^2 R_n| \le \bar{L}_1 \frac{1}{\sqrt{n}},$$

where

$$\bar{L}_1 := L_1 + K_{\Phi 1} + \left(|I_{\Phi 1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi 3}}{3}\right)\frac{1}{N^{1/2}} + K_{\Phi 2}\frac{1}{N}.$$

## Proof of Proposition 2

Fix $c_1^{\max} > c_1$ and $c_1^{\min} \in (\frac{2}{3\sqrt{2}-2}c_1, c_1)$. Let us choose $\delta \in (0,1)$ such that, for $x \in [1-\delta, 1)$,

$$\frac{c_1^{\min}}{(1-x)^{1/2}} \le F_2(x) \le \frac{c_1^{\max}}{(1-x)^{1/2}}, \qquad \frac{1}{2}\frac{c_1^{\min}}{(1-x)^{3/2}} \le F_2'(x) \le \frac{1}{2}\frac{c_1^{\max}}{(1-x)^{3/2}}. \tag{12}$$

Using the representation (1) and (12), we choose $C = C(F) > 0$ such that

$$\int_{-1}^{\beta} |F_2'(x)|dx \le F(\beta) + C, \quad \beta \in [1-\delta, 1). \tag{13}$$

Finally, we assume that (increasing $L_2$ and $l_2$ if necessarily)

$$n \ge \frac{2}{\delta}. \tag{14}$$

Integrating by parts, we have

$$\int_{-1}^{1} F_2(x)dx = \sum_{k=-n}^{n-1} \int_{k/n}^{(k+1)/n} F_2(x)dx$$

$$= \sum_{k=-n}^{n-1} \left( \left( x - \frac{k+1}{n} \right) F_2(x) \Big|_{k/n}^{(k+1)/n} - \int_{k/n}^{(k+1)/n} \left( x - \frac{k+1}{n} \right) F_2'(x)dx \right) = \sum_{k=-n}^{n-1} \frac{1}{n} F_2\left(\frac{k}{n}\right) + \Sigma_{1,n},$$

where

$$\Sigma_{1,n} := \sum_{k=-n}^{n-1} \int_{k/n}^{(k+1)/n} \left( \frac{k+1}{n} - x \right) F_2'(x)dx.$$

Inequalities (12)–(14) imply that

$$\begin{aligned} |\Sigma_{1,n}| &= \left| \int_{1-1/n}^{1} (1-x)F_2'(x) + \sum_{k=-n}^{n-2} \int_{k/n}^{(k+1)/n} \left( \frac{k+1}{n} - x \right) F_2'(x)dx \right| \\ &\le \int_{1-1/n}^{1} \frac{1}{2} \frac{c_1^{\max}}{(1-x)^{1/2}} + \sum_{k=-n}^{n-2} \frac{1}{n} \int_{k/n}^{(k+1)/n} |F_2'(x)|dx \le \frac{c_1^{\max}}{\sqrt{n}} + \frac{1}{n} \int_{-1}^{1-1/n} |F_2'(x)|dx. \end{aligned} \tag{15}$$

Since $1 - 1/n > 1 - \delta$ due to (14), we conclude from (12), (13) and (15) that

$$|\Sigma_{1,n}| \le \frac{c_1^{\max}}{\sqrt{n}} + \frac{1}{n} \left( F_2\left(1 - \frac{1}{n}\right) + C \right) \le 2c_1^{\max} \frac{1}{\sqrt{n}} + C\frac{1}{n}.$$

Taking $L_2 := 2c_1^{\max} + \frac{C}{\sqrt{N}}$, we obtain the second inequality in (2).

Set $k_0 := [(1-\delta)n]$ ($[\cdot]$ is the integer part of a real number). We represent $\Sigma_{1,n}$ as follows:

$$\Sigma_{1,n} = \Sigma_{2,n} + \Sigma_{3,n}, \tag{16}$$

where

$$\begin{aligned} \Sigma_{2,n} &:= \left( \sum_{k=-n}^{k_0-1} \int_{k/n}^{(k+1)/n} \left( \frac{k+1}{n} - x \right) F_2'(x)dx + \int_{k_0}^{1-\delta} \left( \frac{k_0+1}{n} - x \right) F_2'(x)dx \right), \\ \Sigma_{3,n} &:= \left( \int_{1-\delta}^{(k_0+1)/n} \left( \frac{k_0+1}{n} - x \right) F_2'(x)dx + \sum_{k=k_0+1}^{n-1} \int_{k/n}^{(k+1)/n} \left( \frac{k+1}{n} - x \right) F_2'(x) \right). \end{aligned} \tag{17}$$

Below we estimate $\Sigma_{2,n}$ and $\Sigma_{3,n}$ separately. Inequality (13) implies that

$$\Sigma_{2,n} \geq -\frac{1}{n}\left(\int_{-1}^{1-\delta} |F_2'(x)|dx\right) \geq -\frac{1}{n}(F_2(1-\delta)+C). \tag{18}$$

Due to (12) all the terms in expression $\Sigma_{3,n}$ are positive. Inequality (14) implies that the last sum in formula (17) contains at least two terms and inequality (12) implies that

$$\begin{aligned}
\Sigma_{3,n} &\geq \int_{1-2/n}^{1-1/n} \left(1-\frac{1}{n}-x\right) F_2'(x)dx + \int_{1-1/n}^{1} (1-x)F_2'(x)dx \\
&\geq \int_{1-2/n}^{1-1/n} \frac{1}{2}\left(1-\frac{1}{n}-x\right) \frac{c_1^{\min}}{(1-x)^{3/2}}dx + \int_{1-1/n}^{1} \frac{1}{2}\frac{c_1^{\min}}{(1-x)^{1/2}}dx \\
&= \int_{1-2/n}^{1} \frac{1}{2}\frac{c_1^{\min}}{(1-x)^{1/2}}dx - \frac{1}{2n}\int_{1-2/n}^{1-1/n} \frac{c_1^{\min}}{(1-x)^{3/2}} = \frac{3\sqrt{2}-2}{2}c_1^{\min}\frac{1}{\sqrt{n}}.
\end{aligned} \tag{19}$$

Relations (16), (18), and (19) imply that

$$\Sigma_{1,n} \geq L_2^*\frac{1}{\sqrt{n}} - l_2\frac{1}{n},$$

where $L_2^* := \frac{3\sqrt{2}-2}{2}c_1^{\min} > c_1$ and $l_2 := F_2(1-\delta)+C$.

# References


[1] Pavel Gurevich, Sergey Tikhomirov, Spatially discrete reaction-diffusion equations with discontinuous hysteresis, arXiv:1504.02385